arXiv:0912.4101v1 [math.DG] 21 Dec 2009

Bull. Math. Soc. Sc. Math. Roumanie
Tome 39 (87) Nr. 1-4, 1996

Submanifolds of a complex space form, whose geodesics lie in 1 - dimensional complex submanifolds

by

Ognian KASSABOV

In [2], [3] K. Nomizu has proved, that if M^n is a Kaehler submanifold of $P^m(\mathbb{C})$, such that each geodesic of M^n lies in a complex projective line $P^1(\mathbb{C})$ of $P^m(\mathbb{C})$, then M^n is totally geodesic. In this note we generalize this result to an arbitrary submanifold of a complex space form. Namely we prove:

Theorem 1. *Let* M ($\dim_{\mathbb{R}} M \geq 2$) *be a connected, complete submanifold of a complex space form* $\widetilde{M}^m(\mu)$, $\mu \neq 0$. *If each geodesic in* M *lies in an 1-dimensional totally geodesic complex submanifold of* $\widetilde{M}^m(\mu)$, *then* M *is a real or a complex space form and* M *is totally geodesic in* $\widetilde{M}^m(\mu)$.

1 Preliminaries.

Let $\widetilde{M}^m$ be a Kaehler manifold of complex dimension m. Denote by g its metric tensor, by J its complex structure and by $\widetilde{\nabla}$ the covariant differentiation with respect to the Riemannian connection. Then $\widetilde{\nabla} g = 0$ and $\widetilde{\nabla} J = 0$. Let $\widetilde{R}$ be the curvature tensor of $\widetilde{M}^m$. Then $\widetilde{M}^m$ is said to be of constant holomorphic sectional curvature, if there exists a constant μ, such that $\widetilde{R}(x, Jx, Jx, x) = \mu$ for any unit vector x on $\widetilde{M}$. A connected, simply connected, complete Kaehler manifold of constant holomorphic sectional curvature is said to be a complex space form.

Now let M be a submanifold of $\widetilde{M}^m$ and denote by ∇ the covariant differentiation of M with respect to the induced Riemannian connection. Then we write the Gauss formula

$$\widetilde{\nabla}_X Y = \nabla_X Y + \sigma(X,Y) \ ,$$

where σ is a normal-bundle-valued symmetric tensor field on M, called the second fundamental form of M in $\widetilde{M}^m$. We recall that M is called a totally geodesic submanifold of $\widetilde{M}^m$, if $\sigma = 0$. Let ξ be a normal vector field. Then the Weingarten formula is given by

$$\widetilde{\nabla}_X \xi = -A_\xi X + D_X \xi \ ,$$

where $-A_\xi X$ (resp. $D_X \xi$) denotes the tangential (resp. the normal) component of $\widetilde{\nabla}_X \xi$. It is well known that $g(\sigma(X,Y),\xi) = g(A_\xi X, Y)$ and D is the covariant differentiation in the normal bundle.

The submanifold M is said to be a complex (resp. a totally real) submanifold of $\widetilde{M}^m$ if $JT_pM = T_pM$ (resp. $JT_pM = (T_pM)^\perp$) for each point $p \in M$, where T_pM is the tangent space of M at p.

Denote by R the curvature tensor of the Riemannian manifold M. Then M is said to be of constant sectional curvature, if there exists a constant μ, such that $R(x,y,y,x) = \mu$ for any pair $\{x,y\}$ of orthonormal vectors on M. A connected, simply connected, complete Riemannian manifold of constant sectional curvature is called a real space form.

2 Proof of the Theorem.

Let $c = c(t)$ be a geodesic in M, parameterized by arc lenght and denote c' by T. Then

$$\widetilde{\nabla}_T T = \sigma(T,T) \ . \tag{1}$$

By assumption c lies in a 1-dimensional complex totally geodesic submanifold $N^1(\nu)$ of $\widetilde{M}^m(\mu)$. Denote ∇' the covariant differentiation of $N^1(\nu)$. Then $\widetilde{\nabla}_T T = \nabla'_T T = aT + bJT$, where $a = a(t)$, $b = b(t)$. Since c is a geodesic in M $\sigma(T,T) = aT + bJT$ holds good. This implies $ag(T,T) = g(\sigma(T,T),T) = 0$, so $a = 0$ and

$$\sigma(T,T) = bJT \ . \tag{2}$$

Using (1), (2) and $\widetilde{\nabla} J = 0$ we obtain

$$\widetilde{\nabla}_T \sigma(T,T) = T(b)JT - b^2 T \ . \tag{3}$$

On the other hand, according to the Weingarten formula we can write

$$\widetilde{\nabla}_T\sigma(T,T) = -A_{\sigma(T,T)}T + D_T\sigma(T,T)\ .$$

Hence, using (2) and (3) we find easily $A_{\sigma(T,T)}T = b^2T$ and consequently $g(\sigma(x,x),\sigma(x,y)) = 0$ for any orthogonal vectors $x, y \in T_pM$ and any point p of M, so M is an isotropic submanifold of $\widetilde{M}^m(\mu)$, see [4]. Then there exists a function $\lambda(p)$ on M, such that $\lambda(p) \geq 0$ and

$$g(\sigma(x,x),\sigma(x,x)) = \lambda^2(p) \tag{4}$$

for any unit vector $x \in T_pM$ and hence it follows by a standard way

$$g(\sigma(x,x),\sigma(x,x)) + 2g(\sigma(x,y),\sigma(x,y)) = \lambda^2(p) \tag{5}$$

for any orthogonal unit vectors $x, y \in T_pM$, $p \in M$. According to (2), (5) implies

$$g(\sigma(x,y),\sigma(x,y)) = \lambda^2(p) \tag{6}$$

for any orthonormal vectors $x, y \in T_pM$, $p \in M$.

From (2) and (4) it follows

$$\sigma(x,x) = \pm\lambda(p)Jx \tag{7}$$

for any unit vector $x \in T_pM$. Let x, y be orthonormal vectors in T_pM, such that $\sigma(x,x) = \lambda(p)Jx$, $\sigma(y,y) = \lambda(p)Jy$. Then

$$2\sigma(x,y) = \sigma(x+y,x+y) - \sigma(x,x) - \sigma(y,y)$$

and hence, using (7) we obtain

$$\sigma(x,y) = \frac{\epsilon\sqrt{2}-1}{2}\lambda(p)(Jx+Jy)\ ,$$

where $\epsilon = \pm 1$, and consequently

$$g(\sigma(x,y),\sigma(x,y)) = \frac{3-2\epsilon\sqrt{2}}{2}\lambda^2(p)\ . \tag{8}$$

From (6),)7), (8) we conclude that $\lambda = 0$, so M is totally geodesic. Then, according to Theorem 1 in [1] M is a totally real or a complex submanifold of $\widetilde{M}^m(\mu)$, which proves our Theorem.

References

[1] B. Y. CHEN and K. OGIUE : Two theorems on Kaehler manifolds, *Michigan Math. J.*, **21** (1974), 225-229.

[2] K. NOMIZU : Eine Kennzeichnung der projectiven Unterraine in dem Komplexen Projectiven Raum nain ihrem Geodätischen, *Math. Z.*, **137** (1974), 147-150.

[3] K. NOMIZU : A characterization of the Veronese varieties, *Nagoya Math. J.*, **60** (1976), 181-188.

[4] B. O'NEILL : Isotropic and Kaehler immertions, *Canad. J. Math.*, **17** (1965), 905-915.

Higher Transport School
BBTY"T. Kableschkov"
Section of Mathematics
Slatina, 1574 Sofia
BULGARIA

Received September 1, 1995